

Boundary integral equations on unbounded rough surfaces: Fredholmness and the finite section method

SIMON N. CHANDLER-WILDE and MARKO LINDNER

ABSTRACT. We consider a class of boundary integral equations that arise in the study of strongly elliptic BVPs in unbounded domains of the form $D = \{(x, z) \in \mathbb{R}^{n+1} : x \in \mathbb{R}^n, z > f(x)\}$ where $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a sufficiently smooth bounded and continuous function. A number of specific problems of this type, for example acoustic scattering problems, problems involving elastic waves, and problems in potential theory, have been reformulated as second kind integral equations $u + Ku = v$ in the space BC of bounded, continuous functions. Having recourse to the so-called limit operator method, we address two questions for the operator $A = I + K$ under consideration, with an emphasis on the function space setting BC. Firstly, under which conditions is A a Fredholm operator, and, secondly, when is the finite section method applicable to A ?

This project is funded by a Fellowship of the EU (MEIF-CT-2005-009758), which supports the second author.

Authors' address: Department of Mathematics, University of Reading, Whiteknights, PO Box 220, Reading, RG6 6AX, UK.

email: s.n.chandler-wilde@reading.ac.uk and m.lindner@reading.ac.uk

1 Introduction

The boundary integral equation method is very well developed as a tool for the analysis and numerical solution of strongly elliptic boundary value problems in both bounded and unbounded domains, provided the boundary itself is bounded (e.g. [6, 27, 35]).

In the case when both domain and boundary are unbounded, the theory of the boundary integral equation method is much less well developed. The reason for this is fairly clear, namely that loss of compactness of the boundary leads to loss of compactness of boundary integral operators. To be more precise, classical applications of the boundary integral method, for example to potential theory in smooth bounded domains, lead to second kind boundary integral equations of the form $Au = v$ where the function v is known, u unknown, and the operator A is a compact perturbation of the identity (e.g. [6]). In more sophisticated applications, to more complex strongly elliptic systems or to piecewise smooth or general Lipschitz domains, compactness arguments continue to play an impor-

tant role. For example a standard method to establish that a boundary integral operator A is Fredholm of index zero is to show a Gårding inequality, i.e. to establish that A , as an operator on some Hilbert space, is a compact perturbation of an elliptic principal part (e.g. [27]). The case when the boundary is unbounded is difficult because this tool of compactness is no longer available.

To compensate for loss of compactness, only a few alternative tools are known. In the case of classical potential theory and some other strongly elliptic systems, invertibility and/or Fredholmness of boundary integral operators can be established via direct a priori bounds, using Rellich-type identities. In the context of boundary integral equation formulations these arguments were first systematically exploited by Jerison and Kenig [18], Verchota [38] and Dahlberg and Kenig [16] (and see [20, 28]). The main objective in these papers is to overcome loss of compactness associated with *non-smoothness* rather than unboundedness of the boundary, but the Rellich identity arguments used are applicable also when the boundary is infinite in extent, notably, and most straightforwardly, when the boundary is the graph of a Lipschitz function. For example, for classical potential theory, invertibility of the operator $A = I + K$, where I is the identity and K the classical double-layer potential operator, can be established when the boundary is the graph of a Lipschitz function, as discussed in [16, 20, 28]. The Rellich-identity estimates establish invertibility of A in the first instance in L^2 , but, by combining these L^2 estimates with additional arguments, the invertibility of A also in L^p for $2 - \epsilon < p < \infty$ can be established [16, 20]. Here ϵ is some positive constant which depends only on the space dimension and the Lipschitz constant of the boundary.

The same methods of argument can be extended to some other elliptic problems and elliptic systems, e.g. [17, 29, 30]. Recently L^2 solvability has also been established for a second kind integral equation formulation on the (unbounded) graph of a bounded Lipschitz function in a case (the Dirichlet problem for the Helmholtz equation with real wave number) when the associated weak formulation of the boundary value problem is non-coercive [7, 36]. (This lack of coercivity is relatively easily dealt with as a compact perturbation when the boundary is Lipschitz and compact (e.g. [37]), but is much more problematic when the boundary is unbounded.)

In this paper we consider the application of another tool which is available for the study of integral equations on unbounded domains, namely *the limit operator method* [32, 33, 23]. The results we obtain are applicable to the boundary integral equation formulation of many strongly elliptic boundary value problems in unbounded domains of the form

$$D = \{(x, z) \in \mathbb{R}^{n+1} : x \in \mathbb{R}^n, z > f(x)\} \quad (1)$$

where $n \geq 1$ and $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a given bounded and continuous function, in short, $f \in BC$, so that the unbounded boundary is the graph of some bounded function. The results we prove are relevant to the case where the boundary is fairly smooth (Lyapunov), that is f is differentiable with a bounded and α -Hölder continuous gradient for some $\alpha \in (0, 1]$; i.e., for some constant $C > 0$,

$|\nabla f(x) - \nabla f(y)| \leq C|x-y|^\alpha$ holds for all $x, y \in \mathbb{R}^n$. This restriction to relatively smooth boundaries has the implication, for many boundary integral operators on ∂D , for example the classical double-layer potential operator (see §2 below), that loss of compactness arises from the unboundedness of ∂D rather than its lack of smoothness. To be precise, the boundary integral operators we consider, while not compact are nevertheless *locally compact* (in the sense of §3.1), and this local compactness will play a key role in the results we obtain. Throughout we let

$$f_+ = \sup_{x \in \mathbb{R}^n} f(x) \quad \text{and} \quad f_- = \inf_{x \in \mathbb{R}^n} f(x)$$

denote the highest and the lowest elevation of the infinite boundary ∂D . It is convenient to assume, without loss of generality, that $f_- > 0$, so that D is entirely contained in the half space $H = \{(x, z) \in \mathbb{R}^{n+1} : x \in \mathbb{R}^n, z > 0\}$.

Let us introduce the particular class of second kind integral equations on \mathbb{R}^n that we consider in this paper. As we will make clear through detailed examples in §2, equations of this type arise naturally when many strongly elliptic boundary value problems in the domain D are reformulated as boundary integral equations on ∂D . To be specific, boundary value problems arising in acoustic scattering problems [8, 9, 11, 14], in the scattering of elastic waves [2, 3], and in the study of unsteady water waves [31], have all been reformulated as second kind boundary integral equations which, after the obvious parametrization, can be written as

$$u + Ku = v,$$

where K is the integral operator

$$(Ku)(x) = \int_{\mathbb{R}^n} k(x, y) u(y) dy, \quad x \in \mathbb{R}^n \quad (2)$$

with kernel k . Further, in all the above examples, the kernel k has the following particular structure which will be the focus of our study, that

$$k(x, y) = \sum_{i=1}^j b_i(x) k_i(x - y, f(x), f(y)) c_i(y), \quad (3)$$

where

$$b_i \in \text{BC}, \quad k_i \in C((\mathbb{R}^n \setminus \{0\}) \times [f_-, f_+]^2) \quad \text{and} \quad c_i \in L^\infty \quad (4)$$

for $i = 1, \dots, j$, and

$$|k(x, y)| \leq \kappa(x - y), \quad x, y \in \mathbb{R}^n, \quad (5)$$

for some $\kappa \in L^1$. Here and throughout L^p is our abbreviation for $L^p(\mathbb{R}^n)$, for $1 \leq p \leq \infty$, and we denote the norm on L^p by $\|\cdot\|_p$. By $L(L^p)$ and $L(\text{BC})$ we will denote the Banach space of bounded linear operators on L^p and on BC ,

respectively. We note that (2)–(5) imply that $K \in L(L^p)$ for $1 \leq p \leq \infty$ with $\|K\|_{L(L^p)} \leq \|\kappa\|_1$. In particular,

$$\|K\|_{L(L^\infty)} = \sup_{x \in \mathbb{R}^n} \int_{\mathbb{R}^n} |k(x, y)| dy \leq \|\kappa\|_1,$$

and we note that $Ku \in \text{BC}$ for $u \in L^\infty$.

In the cases cited above (see §2 below), the structure (3)–(4) is a simple consequence of the invariance with respect to translations in the plane \mathbb{R}^n of the fundamental solutions used in the integral equation formulations. This property follows in turn from invariance in the \mathbb{R}^n plane of the coefficients in the differential operator. In each case the bound (5) follows from the Hölder continuity of f , which ensures that $k(x, y)$ is only weakly singular at $x = y$, and from the particular choice of fundamental solution used in the integral equation formulation (a Green’s function for the half-plane H in each case), which ensures that $k(x, y)$ decreases sufficiently rapidly as $|x - y| \rightarrow \infty$. Throughout, we will denote the set of all operators K satisfying (2)–(5) for a particular function $f \in \text{BC}$ (but any choices of the functions b_i, k_i, c_i and κ) by \mathcal{K}_f .

There are two main aims of this paper. The major aim is to apply results from the so-called limit operator method [32, 33, 23, 26] to operators satisfying (2)–(5), to address, at least partially, two questions for the operator $A = I + K$:

Fredholmness and Invertibility. Under what conditions is the operator A invertible? More generally, under what conditions is the operator A Fredholm; that is, $Au = 0$ has a finite-dimensional solution space only, and the range of A is closed and has finite co-dimension? So, if A is Fredholm, then the equation $Au = v$ is solvable for all v in a closed subspace of finite co-dimension, and the solution u is unique up to perturbations in a finite-dimensional space.

Applicability of the Finite Section Method. If A is invertible, under which conditions is it possible to replace the equation $Au = v$, i.e.

$$u(x) + \int_{\mathbb{R}^n} k(x, y) u(y) dy = v(x), \quad x \in \mathbb{R}^n, \quad (6)$$

by the finite truncations

$$u_\tau(x) + \int_{|y| \leq \tau} k(x, y) u_\tau(y) dy = v(x), \quad x \in \mathbb{R}^n, \quad (7)$$

with a large $\tau > 0$? In the case when we study equations (6) and (7) in the function space $X = L^\infty$ or $X = \text{BC}$, we say that the method of replacing (6) by (7) is *applicable* if the latter equations are uniquely solvable for all sufficiently large τ and their solutions u_τ converge *strictly* (which means uniformly on every compact set) to the solution u of the original problem (6), for every right-hand side $v \in X$. If this is the case, then we can approximately solve a boundary integral equation on the unbounded surface ∂D by instead solving a boundary integral equation on a large finite truncation of ∂D .

The second aim of the paper, of interest in its own right and helpful to the aim of applying known limit operator results, is to relate operators in the class \mathcal{K}_f , for some $f \in \text{BC}$, to classes of integral operators that have been studied previously in the literature.

Throughout the paper, although many of our results can be extended to other function spaces, especially to L^p for $1 \leq p \leq \infty$, we will concentrate on the case where we view A as an operator on BC . In part we make this restriction just for brevity. Our other reasons for this focus are that, while the function space BC has been the main setting of many of the application-related papers already cited above [8, 9, 11, 14, 2, 3, 31], little has been said in the limit operator literature about Fredholmness and the finite section method in the space BC ; indeed, only recently has the L^p setting for the limiting cases $p = 1, \infty$ been addressed [23, 22, 24, 25, 26]. For $1 < p < \infty$, the Fredholmness of operators $I + K$ on L^p with K locally compact was meanwhile studied using limit operator techniques [34]. We will prove some new results in §3.2 relating, for very general classes of operators, the applicability of the finite section method on BC to its applicability on L^∞ .

The structure and main results of the paper are as follows. In §2 we consider three examples of strongly elliptic boundary value problems in the domain D and exhibit the structure (3)-(5). In §3 we introduce limit operators and related concepts, then in §3.1 we recall recently established sufficient criteria for Fredholmness and necessary conditions for invertibility of an operator A on L^∞ and BC . These criteria, expressed in terms of invertibility of limit operators of the operator A , apply to large classes of operators, but in particular to operators of the class \mathcal{K}_f . In §3.2 we make a preliminary study of the finite section method in the space BC , showing that it is applicable if and only if it is stable and that it is stable on BC if and only if it is stable on L^∞ .

To apply the results of section §3.1 to operators of the class \mathcal{K}_f it is necessary to show that the class of operators considered in §3.1 includes \mathcal{K}_f , and to consider the limit operators of operators in \mathcal{K}_f . As a step in this direction and of interest in its own right, we show in §4.1 that the closure of \mathcal{K}_f in $L(\text{BC})$ is a Banach algebra, in fact the Banach algebra generated by particular combinations of multiplication and convolution operators which we identify. This simplifies the study of the limit operators of $K \in \mathcal{K}_f$, since limit operators of multiplication and convolution operators are well understood (see §2). Note that the observation, which is part of our result, that $K \in \mathcal{K}_f$ can be approximated by finite sums of products of convolution and multiplication operators, has been utilised in particular cases as a computational tool for matrix compression and fast matrix-vector multiplication (see [39] and the references therein). In §4.2, using the results of §4.1, we identify explicitly the limit operators of operators of the class \mathcal{K}_f , in the case when the functions f and the functions b_i and c_i in (3) are sufficiently well-behaved (f , b_i and c_i all uniformly continuous will do), in particular showing that each limit operator of $K \in \mathcal{K}_f$ is in $\mathcal{K}_{\bar{f}}$ for some \bar{f} related to the original function f .

Finally, in §5, we put the results of §3 and §4 together with recent results on the finite section method in L^∞ . Our first main result relates invertibility and Fredholmness of $A = I + K$ as an operator on BC to invertibility of the (explicitly identified) limit operators of A , for $K \in K_f$. Our second result, specific to the case $n = 1$, is a necessary and sufficient criterion for applicability of the finite section method in terms of invertibility of the restrictions to half-lines of the limit operators of A . As a specific example we consider the case when f and the coefficients b_i and c_i are slowly oscillating at infinity when these criteria become very explicit. We also apply our results to the first example of §2 (a boundary integral equation for the Dirichlet problem for the Laplace equation in a non-locally perturbed half-plane).

We finish this introduction by noting that there exist tools which are related to the limit operator method which have been developed by the first author and his collaborators for studying invertibility and the stability and convergence of approximation methods for integral equations on unbounded domains (see [12, 4, 13] and the references therein). These methods can be and have been applied to boundary integral equations of the class that we consider in this paper [8, 9, 11, 14, 2, 3, 31]. We note, however, that no systematic study of operators of the class \mathcal{K}_f has been made in these papers. Moreover, the results in these papers are complementary to those we exhibit here: in particular they lead to sufficient but not necessary conditions for invertibility and applicability of the finite section method, and do not provide criteria for Fredholmness.

2 Examples

We start with some concrete physical problems that have been modelled as elliptic boundary value problem and reformulated as second kind boundary integral equations, the integral operator in each case exhibiting the structure (2)-(5).

Example 2.1. – Potential Theory. In [31] PRESTON, CHAMBERLAIN and CHANDLER-WILDE consider the two-dimensional Dirichlet boundary value problem: Given $\varphi_0 \in BC(\partial D)$, find $\varphi \in C^2(D) \cap BC(\overline{D})$ such that

$$\begin{aligned} \Delta \varphi &= 0 & \text{in } D, \\ \varphi &= \varphi_0 & \text{on } \partial D, \end{aligned}$$

which arises in the theory of classical free surface water wave problems. In this case $n = 1$ and the authors suppose that f is differentiable with bounded and α -Hölder continuous first derivative for some $\alpha \in (0, 1]$, i.e., for some constant $C > 0$, $|f'(x) - f'(y)| \leq C|x - y|^\alpha$ for $x, y \in \mathbb{R}$.

Now let

$$G(\mathbf{x}, \mathbf{y}) = \Phi(\mathbf{x}, \mathbf{y}) - \Phi(\mathbf{x}^r, \mathbf{y}), \quad \mathbf{x}, \mathbf{y} \in \mathbb{R}^2,$$

denote the Green's function for the half plane H where

$$\Phi(\mathbf{x}, \mathbf{y}) = -\frac{1}{2\pi} \ln |\mathbf{x} - \mathbf{y}|_2, \quad \mathbf{x}, \mathbf{y} \in \mathbb{R}^2,$$

with $|\cdot|_2$ denoting the Euclidean norm in \mathbb{R}^2 , is the standard fundamental solution for Laplace's equation in two dimensions, and $\mathbf{x}^r = (x_1, -x_2)$ is the reflection of $\mathbf{x} = (x_1, x_2)$ with respect to ∂H . For the solution of the above boundary value problem the following double layer potential ansatz is made in [31]:

$$\varphi(\mathbf{x}) = \int_{\partial D} \frac{\partial G(\mathbf{x}, \mathbf{y})}{\partial \mathbf{n}(\mathbf{y})} \tilde{u}(\mathbf{y}) ds(\mathbf{y}), \quad \mathbf{x} \in D,$$

where $\mathbf{n}(\mathbf{y}) = (f'(y), -1)$ is a vector normal to ∂D at $\mathbf{y} = (y, f(y))$, and the density function $\tilde{u} \in \text{BC}(\partial D)$ is to be determined. In [31] it is shown that φ satisfies the above Dirichlet boundary value problem if and only if

$$(I - K)\tilde{u} = -2\varphi_0, \quad (8)$$

where

$$(K\tilde{u})(\mathbf{x}) = 2 \int_{\partial D} \frac{\partial G(\mathbf{x}, \mathbf{y})}{\partial \mathbf{n}(\mathbf{y})} \tilde{u}(\mathbf{y}) ds(\mathbf{y}), \quad \mathbf{x} \in \partial D. \quad (9)$$

In accordance with the parametrization $\mathbf{x} = (x, f(x))$ of ∂D , we define

$$u(x) := \tilde{u}(\mathbf{x}) \quad \text{and} \quad b(x) := -2\varphi_0(\mathbf{x}), \quad x \in \mathbb{R},$$

and rewrite equation (8) as the equation

$$u(x) - \int_{-\infty}^{+\infty} k(x, y) u(y) dy = b(x), \quad x \in \mathbb{R}, \quad (10)$$

on the real axis for the unknown function $u \in \text{BC}(\mathbb{R})$, where

$$\begin{aligned} k(x, y) &= 2 \frac{\partial G(\mathbf{x}, \mathbf{y})}{\partial \mathbf{n}(\mathbf{y})} \sqrt{1 + f'(y)^2} = -\frac{1}{\pi} \left(\frac{(\mathbf{x} - \mathbf{y}) \cdot \mathbf{n}(\mathbf{y})}{|\mathbf{x} - \mathbf{y}|_2^2} - \frac{(\mathbf{x}^r - \mathbf{y}) \cdot \mathbf{n}(\mathbf{y})}{|\mathbf{x}^r - \mathbf{y}|_2^2} \right) \\ &= -\frac{1}{\pi} \left(\frac{(x - y)f'(y) - f(x) + f(y)}{(x - y)^2 + (f(x) - f(y))^2} - \frac{(x - y)f'(y) + f(x) + f(y)}{(x - y)^2 + (-f(x) - f(y))^2} \right) \\ &= -\frac{1}{\pi} \left(\frac{x - y}{(x - y)^2 + (f(x) - f(y))^2} - \frac{x - y}{(x - y)^2 + (f(x) + f(y))^2} \right) f'(y) \\ &\quad + \frac{1}{\pi} \left(\frac{f(x) - f(y)}{(x - y)^2 + (f(x) - f(y))^2} + \frac{f(x) + f(y)}{(x - y)^2 + (f(x) + f(y))^2} \right). \end{aligned}$$

Clearly $k(x, y)$ is of the form (3) with $j = 2$ and property (4) satisfied. From Lemma 2.1 and inequality (5) in [31] we moreover get that the inequality (5) holds with

$$\kappa(x) = \begin{cases} c|x|^{\alpha-1} & \text{if } 0 < |x| \leq 1, \\ c|x|^{-2} & \text{if } |x| > 1, \end{cases}$$

where $\alpha \in (0, 1]$ is the Hölder exponent of f' , and c is some positive constant. \square

Example 2.2. – Wave scattering by an unbounded rough surface. In [9] CHANDLER-WILDE, ROSS and ZHANG consider the corresponding problem for the Helmholtz equation in two dimensions. Given $\varphi_0 \in \text{BC}(\partial D)$, they seek $\varphi \in C^2(D) \cap \text{BC}(\overline{D})$ such that

$$\begin{aligned} \Delta\varphi + k^2\varphi &= 0 & \text{in } D, \\ \varphi &= \varphi_0 & \text{on } \partial D, \end{aligned}$$

and such that φ satisfies an appropriate radiation condition and constraints on growth at infinity. Again, $n = 1$ and the surface function f is assumed to be differentiable with a bounded and α -Hölder continuous first derivative for some $\alpha \in (0, 1]$. This problem models the scattering of acoustic waves by a sound-soft rough surface; the same problem arises in time-harmonic electromagnetic scattering by a perfectly conducting rough surface.

The authors reformulate this problem as a boundary integral equation which has exactly the form (8)-(9), except that $G(\mathbf{x}, \mathbf{y})$ is now defined to be the Green's function for the Helmholtz equation in the half-plane H which satisfies the impedance condition $\partial G/\partial x_2 + ikG = 0$ on ∂H . As in Example 2.1, this boundary integral equation can be written in the form (10) with $k(x, y)$ of the form (3) with $j = 2$ and property (4) satisfied, and also here inequality (5) holds with

$$\kappa(x) = \begin{cases} c|x|^{\alpha-1} & \text{if } 0 < |x| \leq 1, \\ c|x|^{-3/2} & \text{if } |x| > 1, \end{cases}$$

where $\alpha \in (0, 1]$ is the Hölder exponent of f' , and c is some positive constant. \square

Example 2.3. – Wave propagation over a flat inhomogeneous surface. The propagation of mono-frequency acoustic or electromagnetic waves over flat inhomogeneous terrain has been modelled in two dimensions by the Helmholtz equation

$$\Delta\varphi + k^2\varphi = 0$$

in the upper half plane $D = H$ (so $f \equiv 0$ in (1)) with a Robin (or impedance) condition

$$\frac{\partial\varphi}{\partial x_2} + ik\beta\varphi = \varphi_0$$

on the boundary line ∂D . Here k , the wavenumber, is constant, $\beta \in L^\infty(\partial D)$ is the surface admittance describing the local properties of the ground surface ∂D , and the inhomogeneous term φ_0 is in $L^\infty(\partial D)$ as well.

Similarly to Example 2.2, in fact using the same Green's function $G(\mathbf{x}, \mathbf{y})$ for the Helmholtz equation, CHANDLER-WILDE, RAHMAN and ROSS [11] reformulate this problem as a boundary integral equation on the real line,

$$u(x) - \int_{-\infty}^{+\infty} \tilde{\kappa}(x-y)z(y)u(y) dy = \psi(x), \quad x \in \mathbb{R}, \quad (11)$$

where $\psi \in \text{BC}$ is given and $u \in \text{BC}$ is to be determined. The function $\tilde{\kappa}$ is in $L^1 \cap C(\mathbb{R} \setminus \{0\})$, and $z \in L^\infty$ is closely connected with the surface admittance β by $z = i(1 - \beta)$.

Note that the kernel function of the integral operator in (11) is of the form (3) with $j = 1$. The validity of (4) and (5) is trivial in this case. \square

3 Limit Operators and Finite Sections

The key to both the Fredholm property and the applicability of the finite section method is the behaviour of our operator A at infinity. The tool we shall use to study this behaviour is the so-called limit operator method.

3.1 Limit Operators

Roughly speaking, a limit operator of A is a local representative of A at infinity – a possibly simpler operator that reflects how A acts out there. For its definition we need the following preliminaries.

For every $h \in \mathbb{R}^n$, let V_h denote the *shift operator* acting on $L^p(\mathbb{R}^n)$ by $(V_h u)(x) = u(x - h)$ for all $x \in \mathbb{R}^n$. For every measurable set $U \subset \mathbb{R}^n$, let P_U refer to the operator of multiplication by the characteristic function of U , and write P_τ for P_U if U is the ball around the origin with radius $\tau > 0$. We say that a sequence $(f_k) \subset L^\infty$ *converges strictly* to $f \in L^\infty$ if $\sup \|f_k\|_\infty < \infty$ and $\|P_m(f_k - f)\|_\infty \rightarrow 0$ for all $m \in \mathbb{N}$ as $k \rightarrow \infty$. Finally, a sequence of bounded linear operators (A_k) on L^p is said to \mathcal{P} -*converge* to A if $\sup \|A_k\| < \infty$ and $\|P_m(A_k - A)\|$ and $\|(A_k - A)P_m\|$ tend to zero for all $m \in \mathbb{N}$ as $k \rightarrow \infty$.

Definition 3.1 *If $p \in [1, \infty]$, A is a bounded linear operator on L^p and $h = (h_1, h_2, \dots) \subset \mathbb{Z}^n$ is a sequence tending to infinity, then the \mathcal{P} -limit of the sequence $V_{-h_k} A V_{h_k}$, if it exists, is called the limit operator of A with respect to h and it is denoted by A_h .*

Example 3.2. As a simple example, if $A = M_b$ is the operator of multiplication by the function $b \in L^\infty$, and if $h = (h_1, h_2, \dots)$ tends to infinity, then the limit operator A_h exists if and only if the sequence $V_{-h_k} b = b(\cdot + h_k)$ strictly converges to a function, say $b^{(h)}$, as $k \rightarrow \infty$, in which case $A_h = M_{b^{(h)}}$ is the operator of multiplication by $b^{(h)}$. \square

Example 3.3. The limit operators of M_b are particularly simple if b is what we call a slowly oscillating function. A function $b \in L^\infty$ is *slowly oscillating* if

$$\text{ess sup}_{|y| \leq 1} |b(x + y) - b(x)| \rightarrow 0 \quad \text{as} \quad x \rightarrow \infty.$$

In this case, using the notation of Example 3.2, the strict limit $b^{(h)}$, whenever it exists, is just a constant function in the local essential range of b at infinity,

and conversely, every function of this type is a strict limit $b^{(h)}$ with a suitable sequence $h = (h_1, h_2, \dots) \subset \mathbb{Z}^n$ tending to infinity [25].

If, for example, $n = 1$ and $b(x) = \sin \sqrt{|x|}$ and h_k is the integer part of $k^2 \pi^2$ for $k = 1, 2, \dots$, then $b^{(h)}$ exists and is the zero function. It is easily seen that all limit operators of M_b are of the form cI with a constant $c \in [-1, 1]$, and vice versa. \square

A bounded linear operator A on L^p is *band-dominated* if $\sup \|P_U A P_V\| \rightarrow 0$ as $d \rightarrow \infty$, where the supremum is taken over all measurable $U, V \subset \mathbb{R}^n$ with $\text{dist}(U, V) \geq d$. An operator A on L^p is called *rich* if, from every sequence $h \subset \mathbb{Z}^n$ tending to infinity, we can choose a subsequence g such that the limit operator A_g exists. We note that both the band-dominated operators and the rich operators are Banach subalgebras of $L(L^p)$ (see e.g. [26]). Finally, A is called *locally compact* (on L^p) if $P_\tau A$ and $A P_\tau$ are compact for all $\tau > 0$.

Note that operators in the class \mathcal{K}_f , for some $f \in \text{BC}$, are band dominated and locally compact as operators on L^p for $1 \leq p \leq \infty$. In the case $p = \infty$ this will be shown in §4. In §4.2 we will study the limit operators of operators $K \in \mathcal{K}_f$ and will show that such operators are rich (on L^∞) if f is uniformly continuous and if the operators of multiplication by b_i and c_i (b_i and c_i as in (3)) are rich, which, for example, is the case if each of b_i and c_i is uniformly continuous. We note also that, defining, for each $K \in \mathcal{K}_f$, K^* to be the integral operator defined by (2), with $k(x, y)$ replaced by $k(y, x)$, it follows easily from Fubini's theorem that

$$\int_{\mathbb{R}^n} \phi(K^* \psi) dx = \int_{\mathbb{R}^n} (K \phi) \psi dx,$$

for $\phi \in L^p$, $\psi \in L^q$, $1 \leq p \leq \infty$, where $\frac{1}{p} + \frac{1}{q} = 1$. Since L^q can be identified with the dual of L^p for $1 \leq p < \infty$, the operator $K^* \in L(L^q)$ is the adjoint of $K \in L(L^p)$ for $1 \leq p < \infty$. The case $p = \infty$ is an anomaly here, but we can say that $K^* \in L(L^1)$ is the *pre-adjoint* of $K \in L(L^\infty)$ (which just means that K is the adjoint of K^*). This observation is relevant to the next theorem. Note that, in the case that the functions c_i in the definition (3) of $K \in \mathcal{K}_f$ are continuous, K^* is also in \mathcal{K}_f .

The following theorems are the known results on Fredholmness and invertibility from the theory of limit operators that we will apply in §5 to operators $K \in \mathcal{K}_f$, after studying the limit operators of $K \in \mathcal{K}_f$ in §4. The first theorem, a sufficient condition for Fredholmness, is a rather deep result. The second result, which is much more straightforward, is a necessary condition for invertibility.

Theorem 3.4 *If $A = I + K$ and, as an operator on L^∞ , K is band-dominated, rich and locally compact, then the following holds. If all limit operators of A are invertible on L^∞ , then A is Fredholm as an operator on L^∞ . If also $K(L^\infty) \subset \text{BC}$ then A is also Fredholm if restricted to BC .*

Proof. Let all limit operators of A be invertible on L^∞ . From Theorem 2 in [23] and the *if* part of Theorem 1.1 in [22] (alias Theorem 1 in [23]) it follows that

A is invertible at infinity, as defined in [22, 23]. Note that the *if* portion of this Theorem 1.1 does require a rich operator but not the existence of a pre-adjoint (also see Remark 3.5 in [22]).

This, together with K being locally compact implies that $A = I + K$ is Fredholm on L^∞ , by [15, Subsection 3.3]. Further, if also $K(L^\infty) \subset \text{BC}$, then, by Lemma 3.8 c) below, A is Fredholm if restricted to BC . ■

Note that, for $1 < p < \infty$, the invertibility of all limit operators of A (and the uniform boundedness of their inverses) is even necessary and sufficient for the Fredholmness of $A = I + K$ on L^p (see [34]).

Theorem 3.5 *If, as an operator on L^∞ , A is band-dominated, possesses a pre-adjoint in $L(L^1)$ and is invertible, then all limit operators of A are invertible on L^∞ .*

Proof. This follows from the *only if* portion of Theorem 1.1 in [22]. The operator need not be rich for this implication, as pointed out in Remark 3.5 of [22]. ■

We introduce two types of linear operators which will serve as basic building blocks for the operators we study in the rest of the paper. Firstly, for a function $b \in L^\infty$, let M_b denote the *multiplication operator* $u \mapsto bu$. Secondly, for $\kappa \in L^1$, with Fourier Transform a given by

$$a(\xi) = F\kappa(\xi) = \int_{\mathbb{R}^n} e^{i\xi \cdot y} \kappa(y) dy, \quad \xi \in \mathbb{R}^n,$$

where \cdot is the Euclidean inner product on \mathbb{R}^n , let C_a denote the *operator of convolution with κ* , defined by

$$(C_a u)(x) = \int_{\mathbb{R}^n} \kappa(x - y) u(y) dy, \quad x \in \mathbb{R}^n.$$

Moreover, let $FL^1 = \{F\kappa : \kappa \in L^1\}$. It is well known (e.g. Jörgens [19]) that, for $1 \leq p \leq \infty$, the spectrum of C_a as an element of $L(L^p)$ is $\{a(\xi) : \xi \in \mathbb{R}^n\} \cup \{0\}$.

We will denote the set of all $b \in L^\infty$ for which M_b is a rich operator by $L_{\mathfrak{S}}^\infty$. So, by Example 3.2, $b \in L_{\mathfrak{S}}^\infty$ iff every sequence in \mathbb{Z}^n tending to infinity has an infinite subsequence $h = (h_m)$ such that there exists a function $c \in L^\infty$ with

$$\|b|_{h_m+U} - c|_U\|_\infty \rightarrow 0 \quad \text{as } m \rightarrow \infty \quad (12)$$

for every compact set $U \subset \mathbb{R}^n$. A straightforward computation shows that the operator $C_a M_b$ with $a \in FL^1$ is rich as an operator on L^∞ if the above holds with (12) replaced by the much weaker condition

$$\|b|_{h_m+U} - c|_U\|_1 \rightarrow 0 \quad \text{as } m \rightarrow \infty. \quad (13)$$

We denote the set of all $b \in L^\infty$ with this property by $L_{\mathfrak{S}\text{CS}}^\infty$, and write $\tilde{b}^{(h)}$ for the function c with property (13) for all compact sets U . Recall from Example 3.2 that we write $b^{(h)}$ for the function c in (12).

3.2 The finite section method for BC

In this section we will briefly introduce an approximation method for operators on the space of bounded and continuous functions $BC \subset L^\infty$. The operators of interest to us will be of the form

$$A = I + K,$$

where K shall be bounded and linear on L^∞ with the condition $Ku \in BC$ for all $u \in L^\infty$. Typically, K will be some integral operator. One of the simplest examples is a convolution operator $K = C_a$ with $a \in FL^1$. The following lemma follows easily from the denseness in L^1 of the set of C^∞ compactly supported functions.

Lemma 3.6 *If $a \in FL^1$, then $C_a u$ is a continuous function for every $u \in L^\infty$.*

As a slightly more sophisticated example, one could look at an operator of the following form or at the norm limit of a sequence of such operators.

Example 3.7. Put

$$K := \sum_{i=1}^j M_{b_i} C_{a_i} M_{c_i}, \quad (14)$$

where $b_i \in BC$, $a_i \in FL^1$, $c_i \in L^\infty$ and $j \in \mathbb{N}$. For the condition that K maps L^∞ into BC , it is sufficient to impose continuity of the functions b_i in (14), whereas the functions c_i need not be continuous since their action is smoothed by the convolution thereafter. \square

The first two statements of the following auxiliary result are essentially Lemma 3.4 from [4].

Lemma 3.8 *Suppose that $A = I + K$ and that $K \in L(L^\infty)$ and $K(L^\infty) \subset BC$. Abbreviate the restriction $A|_{BC}$ by A_0 . Then the following hold:*

- a) $Au \in BC$ if and only if $u \in BC$;
- b) A is invertible on L^∞ if and only if A_0 is invertible on BC . In this case

$$\|A_0^{-1}\|_{L(BC)} \leq \|A^{-1}\|_{L(L^\infty)} \leq 1 + \|A_0^{-1}\|_{L(BC)} \|K\|_{L(L^\infty)}. \quad (15)$$

- c) If A is a Fredholm operator on L^∞ , then A_0 is Fredholm on BC .

Proof. a) This is immediate from $Au = u + Ku$ and $Ku \in BC$ for all $u \in L^\infty$.

b) If A is invertible on L^∞ , then the invertibility of A_0 on BC and the first inequality in (15) follows from a).

Now let A_0 be invertible on BC . To see that A is injective on L^∞ , suppose $Au = 0$ for $u \in L^\infty$. From $0 \in BC$ and a) we get that $u \in BC$ and hence $u = 0$

since A is injective on BC . Surjectivity of A on L^∞ : Since A_0 is surjective on BC , for every $v \in L^\infty$ there is a $u \in BC$ such that $A_0u = Kv \in BC$. Consequently,

$$A(v - u) = Av - A_0u = v + Kv - Kv = v \quad (16)$$

holds, showing the surjectivity of A on L^∞ . So A is invertible on L^∞ , and, by (16), $A^{-1}v = v - u = v - A_0^{-1}Kv$ for all $v \in L^\infty$, and hence $A^{-1} = I - A_0^{-1}K$. This proves the second inequality in (15).

c) From a) we get that $\ker A \subset BC$ since $0 \in BC$. But this implies that

$$\ker A_0 = \ker A. \quad (17)$$

Another immediate consequence of a) is

$$A_0(BC) = A(L^\infty) \cap BC. \quad (18)$$

Finally, by (18), we have the following relation between factor spaces

$$\frac{BC}{A_0(BC)} = \frac{BC}{A(L^\infty) \cap BC} \cong \frac{BC + A(L^\infty)}{A(L^\infty)} \subset \frac{L^\infty}{A(L^\infty)}. \quad (19)$$

So, if A is Fredholm on L^∞ , then (17), (19) and (18) show that also $\ker A_0$ and $BC/A_0(BC)$ are finite-dimensional and $A_0(BC)$ is closed. ■

Remark 3.9. a) The previous lemma clearly holds for arbitrary Banach spaces with one of them contained in the other in place of BC and L^∞ .

b) If, moreover, K has a pre-adjoint operator on L^1 , then an approximation argument as in the proof of Lemma 3.5 of [4] even shows that, in fact, (15) can be improved to the equality $\|A_0^{-1}\|_{L(BC)} = \|A^{-1}\|_{L(L^\infty)}$. □

This paper is concerned with the equation $Au = b$ with $A = I + K$, particularly with the case where $u \in L^\infty$ and $b \in BC$ and the equation $Au = b$ is some integral equation

$$u(x) + \int_{\mathbb{R}^n} k(x, y) u(y) dy = b(x), \quad x \in \mathbb{R}^n. \quad (20)$$

In this case, by Lemma 3.8 a), we are looking for u in BC only.

In this setting, a popular approximation method which dates back at least to Atkinson [5] and Anselone and Sloan [1], is just to reduce the range of integration from \mathbb{R}^n to the ball $|y| \leq \tau$ for some $\tau > 0$. We call this procedure the *finite section method* for BC (short: BC -FSM). We are now looking for solutions $u_\tau \in BC$ of

$$u_\tau(x) + \int_{|y| \leq \tau} k(x, y) u_\tau(y) dy = b(x), \quad x \in \mathbb{R}^n \quad (21)$$

with $\tau > 0$, and hope that the sequence (u_τ) of solutions of (21) strictly converges to the solution u of (20) as $\tau \rightarrow \infty$.

This method (21) can be written as $A_\tau u_\tau = b$ with

$$A_\tau = I + KP_\tau. \quad (22)$$

As a consequence of Lemma 3.8 a) applied to A_τ , one also has

Corollary 3.10 *For every $\tau > 0$, it holds that $A_\tau u_\tau \in \text{BC}$ iff $u_\tau \in \text{BC}$.*

We say that a sequence of operators (A_τ) is *stable* if there exists a τ_0 such that all A_τ with $\tau > \tau_0$ are invertible and their inverses are uniformly bounded.

In accordance with the machinery presented in [33, 23], our strategy to study equation (20) and the stability of its approximation by (21) is to embed these into L^∞ , where we can relate the BC-FSM (21) to the usual FSM

$$P_\tau A P_\tau u_\tau = P_\tau b \quad (23)$$

on L^∞ . Indeed, the applicabilities of these different methods turn out to be equivalent.

Proposition 3.11 *For the operator $A = I + K$ with $K(L^\infty) \subset \text{BC}$, let*

$$A_\tau := I + KP_\tau \quad \text{and} \quad A_{[\tau]} := P_\tau A P_\tau + Q_\tau, \quad \tau > 0.$$

Then the following statements are equivalent.

- (i) *The BC-FSM (A_τ) alias (21) is applicable in BC.*
- (ii) *The BC-FSM (A_τ) alias (21) is applicable in L^∞ .*
- (iii) *The FSM $(A_{[\tau]})$ is applicable in L^∞ .*
- (iv) *(A_τ) is stable on BC.*
- (v) *(A_τ) is stable on L^∞ .*
- (vi) *$(A_{[\tau]})$ is stable on L^∞ .*

Proof. The implication (i) \Rightarrow (iv) is standard. The equivalence of (iv) and (v) follows from Lemma 3.8 b) applied to A_τ . The equivalence of (v) and (vi) was already pointed out in [21]. It comes from the following observation:

$$\begin{aligned} A_\tau = I + KP_\tau &= P_\tau + P_\tau K P_\tau + Q_\tau + Q_\tau K P_\tau \\ &= P_\tau A P_\tau + Q_\tau (I + Q_\tau K P_\tau) \\ &= (P_\tau A P_\tau + Q_\tau) (I + Q_\tau K P_\tau) \\ &= A_{[\tau]} (I + Q_\tau K P_\tau), \end{aligned}$$

where the second factor $(I + Q_\tau K P_\tau)$ is always invertible with its inverse equal to $I - Q_\tau K P_\tau$, and hence $\|(I + Q_\tau K P_\tau)^{-1}\| \leq 1 + \|K\|$ for all $\tau > 0$.

(v) \Rightarrow (ii). Since (v) implies (vi), it also implies the invertibility of A on L^∞ by Theorem 4.2 in [24] (Theorem 5.2 in [23]). But this, together with (v), implies (ii) by Theorem 1.44 in [23].

Finally, the implication (ii) \Rightarrow (i) is trivial if we keep in mind Lemma 3.8 a) and Corollary 3.10, and the equivalence of (iii) and (vi) follows from Theorem 4.2 in [24]. ■

For the study of property (iii) in Proposition 3.11 we have Theorem 4.2 in [24] (alias Theorem 5.2 in [23]) involving limit operators of A , provided that, in addition, A is a rich operator. We will state the final result in Section 5.2.

4 Properties of Integral Operators of the Class \mathcal{K}_f

Recall that the aim of the paper is to study the operator $A = I + K$ where K is subject to (2)–(5), and that, for a given function $f \in \text{BC}$, we denote the class of all these operators K by \mathcal{K}_f .

4.1 The relationship between \mathcal{K}_f and other classes of integral operators

For technical reasons we find it convenient to embed this class into a somewhat larger Banach algebra of integral operators. (It will turn out that this Banach algebra actually is not that much larger than \mathcal{K}_f). Therefore, given $f \in \text{BC}$, put $f_- := \inf f$, $f_+ := \sup f$, and let R_f denote the set of all operators of the form

$$(Bu)(x) = \int_{\mathbb{R}^n} k(x-y, f(x), f(y)) u(y) dy, \quad x \in \mathbb{R}^n \quad (24)$$

with $k \in C(\mathbb{R}^n \times [f_-, f_+]^2)$ compactly supported. Moreover, put

$$\begin{aligned} \hat{\mathcal{B}} &:= \text{closspan}\{ M_b B M_c : b \in \text{BC}, B \in R_f, c \in L^\infty \}, \\ \mathcal{B} &:= \text{closalg}\{ M_b B M_c : b \in \text{BC}, B \in R_f, c \in L^\infty \}, \\ \hat{\mathcal{C}} &:= \text{closspan}\{ M_b C_a M_c : b \in \text{BC}, a \in FL^1, c \in L^\infty \}, \\ \mathcal{C} &:= \text{closalg}\{ M_b C_a M_c : b \in \text{BC}, a \in FL^1, c \in L^\infty \}, \\ \mathcal{A} &:= \text{closalg}\{ M_b, C_a : b \in L^\infty, a \in FL^1 \}. \end{aligned}$$

Remark 4.1. a) Here, $\text{closspan } M$ denotes the closure in $L(\text{BC})$ of the set of all finite sums of elements of $M \subset L(\text{BC})$, and $\text{closalg } M$ denotes the closure in $L(\text{BC})$ of the set of all finite sum-products of elements of M . So $\text{closspan } M$ is the smallest closed subspace and $\text{closalg } M$ the smallest (not necessarily unital) Banach subalgebra of $L(\text{BC})$ containing M . In both cases we say they are *generated by* M .

b) The following proposition shows that $\hat{\mathcal{B}}$ and \mathcal{B} do not depend on the function $f \in \text{BC}$ which is why we omit f in their notations.

c) It is easily seen (see Example 3.7) that all operators in $\hat{\mathcal{C}}$ map arbitrary elements from L^∞ into BC . Consequently, every $K \in \hat{\mathcal{C}}$ is subject to the condition on K in Subsection 3.2.

d) The linear space $\hat{\mathcal{C}}$ is the closure of the set of operators considered in Example 3.7. The following proposition shows that this set already contains all of \mathcal{K}_f . More precisely, it coincides with the closure of \mathcal{K}_f in the norm of $L(\text{BC})$ and with the other spaces and algebras introduced above. \square

Proposition 4.2 *The identity*

$$\text{clos } \mathcal{K}_f = \hat{\mathcal{B}} = \hat{\mathcal{C}} = \mathcal{B} = \mathcal{C} \subset \mathcal{A}$$

holds.

Proof. Clearly, $\hat{\mathcal{C}} \subset \hat{\mathcal{B}}$ since C_a with $a \in FL^1$ can be approximated in the operator norm by convolutions $B = C_{a'}$ with a continuous and compactly supported kernel. But these operators B are clearly in R_f .

For the reverse inclusion, $\hat{\mathcal{B}} \subset \hat{\mathcal{C}}$, it is sufficient to show that the generators of $\hat{\mathcal{B}}$ are contained in $\hat{\mathcal{C}}$. We will prove this by showing that $B \in \hat{\mathcal{C}}$ for all $B \in R_f$. So let $k \in C(\mathbb{R}^n \times [f_-, f_+]^2)$ be compactly supported, and define B as in (24). To see that $B \in \hat{\mathcal{C}}$, take $L \in \mathbb{N}$, choose $f_- = s_1 < s_2 < \dots < s_{L-1} < s_L = f_+$ equidistant in $[f_-, f_+]$, and let φ_ξ denote the standard Courant hat function for this mesh that is centered at s_ξ . Then, since k is uniformly continuous, its piecewise linear interpolations (with respect to the variables s and t),

$$k^{(L)}(r, s, t) := \sum_{\xi, \eta=1}^L k(r, s_\xi, s_\eta) \varphi_\xi(s) \varphi_\eta(t), \quad r \in \mathbb{R}^n, \quad s, t \in [f_-, f_+],$$

uniformly approximate k as $L \rightarrow \infty$, whence the corresponding integral operators with k replaced by $k^{(L)}$ in (24),

$$(B^{(L)}u)(x) = \int_{\mathbb{R}^n} \sum_{\xi, \eta=1}^L k(x-y, s_\xi, s_\eta) \varphi_\xi(f(x)) \varphi_\eta(f(y)) u(y) dy, \quad (25)$$

converge to B in the operator norm in $L(\text{BC})$ as $L \rightarrow \infty$. But it is obvious from (25) that $B^{(L)} \in \hat{\mathcal{C}}$, which proves that also $B \in \hat{\mathcal{C}}$.

To see that $\mathcal{B} = \mathcal{C}$, it is sufficient to show that the generators of each of the algebras are contained in the other algebra. But this follows from $\hat{\mathcal{B}} = \hat{\mathcal{C}}$, which is already proven.

That \mathcal{C} is contained in the Banach algebra \mathcal{A} generated by L^1 -convolutions and L^∞ -multiplications, is obvious.

For the inclusion $\mathcal{C} \subset \hat{\mathcal{C}}$ it is sufficient to show that $C_a M_b C_c \in \hat{\mathcal{C}}$ for all $a, c \in FL^1$ and $b \in L^\infty$. So take an arbitrary $b \in L^\infty$ and let $a = F\kappa$ and $c = F\lambda$ with $\kappa, \lambda \in L^1$. Arguing as in the first paragraph of the proof, it is sufficient to consider the case where κ and λ are continuous and compactly supported, say $\kappa(x) = \lambda(x) = 0$ if $|x| > \ell$. It is now easily checked that

$$(C_a M_b C_c u)(x) = \int_{\mathbb{R}^n} k(x, y) u(y) dy, \quad x \in \mathbb{R}^n,$$

with

$$k(x, y) = \int_{\mathbb{R}^n} \kappa(x - z) b(z) \lambda(z - y) dz = \int_{|t| \leq \ell} \kappa(t) b(x - t) \lambda(x - t - y) dt.$$

By taking a sufficiently fine partition into measurable subsets, $\{T_1, \dots, T_N\}$, of $\{t : |t| < \ell\}$, that is a partition with $\max_i \text{diam } T_i$ sufficiently small, and fixing $t_m \in T_m$ for $m = 1, \dots, N$, we can approximate $k(x, y)$ arbitrarily closely in the supremum norm on $\mathbb{R}^n \times \mathbb{R}^n$ by

$$k(x, y) = \sum_{m=1}^N \int_{T_m} \kappa(t) b(x - t) \lambda(x - t - y) dt \approx k_N(x, y)$$

where

$$\begin{aligned} k_N(x, y) &:= \sum_{m=1}^N \kappa(t_m) \lambda(x - t_m - y) \int_{T_m} b(x - t) dt \\ &= \sum_{m=1}^N \kappa_m \lambda_m(x - y) b_m(x), \quad x, y \in \mathbb{R}^n, \end{aligned} \quad (26)$$

with $\kappa_m = \kappa(t_m)$, $\lambda_m(x) = \lambda(x - t_m)$ and $b_m(x) = \int_{T_m} b(x - t) dt$, the latter depending continuously on x . In particular, choosing the partition so that $\max_i \text{diam } T_i < 1/N$, and noting that $k(x, y) = k_N(x, y) = 0$ for $|x - y| > 2\ell$, we see that

$$\sup_{x \in \mathbb{R}^n} \int_{\mathbb{R}^n} |k(x, y) - k_N(x, y)| dy \rightarrow 0 \text{ as } N \rightarrow \infty,$$

so that $C_a M_b C_c \in \hat{\mathcal{C}}$.

The inclusion $\hat{\mathcal{B}} \subset \text{clos } \mathcal{K}_f$ is also obvious since (4) and (5) hold if $b_i \in \text{BC}$, $c_i \in L^\infty$ and k_i is compactly supported and continuous on all of $\mathbb{R}^n \times [f_-, f_+]^2$.

So it remains to show that $\text{clos } \mathcal{K}_f \subset \hat{\mathcal{B}}$. This clearly follows if we show that $\mathcal{K}_f \subset \hat{\mathcal{B}}$. So let $K \in \mathcal{K}_f$ be arbitrary, that means K is an integral operator of the form (2) with a kernel $k(\cdot, \cdot)$ subject to (3), (4) and (5). For every $\ell \in \mathbb{N}$, let $p_\ell : [0, \infty) \rightarrow [0, 1]$ denote a continuous function with support in $[1/(2\ell), 2\ell]$ which is identically equal to 1 on $[1/\ell, \ell]$. Then, for $i = 1, \dots, j$,

$$k_i^{(\ell)}(r, s, t) := p_\ell(|r|) k_i(r, s, t), \quad r \in \mathbb{R}^n, s, t \in [f_-, f_+],$$

is compactly supported and continuous on $\mathbb{R}^n \times [f_-, f_+]^2$, whence $B_i^{(\ell)} \in R_f$ with

$$(B_i^{(\ell)} u)(x) := \int_{\mathbb{R}^n} k_i^{(\ell)}(x - y, f(x), f(y)) u(y) dy, \quad x \in \mathbb{R}^n,$$

for all $u \in \text{BC}$. Now put

$$\begin{aligned} k^{(\ell)}(x, y) &:= \sum_{i=1}^j b_i(x) k_i^{(\ell)}(x-y, f(x), f(y)) c_i(y) \\ &= p_\ell(|x-y|) k(x, y), \end{aligned}$$

and let $K^{(\ell)}$ denote the operator (2) with k replaced by $k^{(\ell)}$; that is

$$K^{(\ell)} = \sum_{i=1}^j M_{b_i} B_i^{(\ell)} M_{c_i}, \quad (27)$$

which is clearly in $\hat{\mathcal{B}}$. It remains to show that $K^{(\ell)} \rightrightarrows K$ as $\ell \rightarrow \infty$. Therefore, note that

$$\begin{aligned} \|K - K^{(\ell)}\| &\leq \sup_{x \in \mathbb{R}^n} \int_{\mathbb{R}^n} |k(x, y) - k^{(\ell)}(x, y)| dy \\ &= \sup_{x \in \mathbb{R}^n} \int_{\mathbb{R}^n} |(1 - p_\ell(|x-y|)) k(x, y)| dy \\ &\leq \sup_{x \in \mathbb{R}^n} \int_{|x-y| < 1/\ell} |k(x, y)| dy + \sup_{x \in \mathbb{R}^n} \int_{|x-y| > \ell} |k(x, y)| dy \\ &\leq \sup_{x \in \mathbb{R}^n} \int_{|x-y| < 1/\ell} |\kappa(x-y)| dy + \sup_{x \in \mathbb{R}^n} \int_{|x-y| > \ell} |\kappa(x-y)| dy \\ &= \int_{|z| < 1/\ell} |\kappa(z)| dz + \int_{|z| > \ell} |\kappa(z)| dz \end{aligned}$$

with $\kappa \in L^1$ from (5). But clearly, this goes to zero as $\ell \rightarrow \infty$. ■

4.2 The limit operators of integral operators in \mathcal{K}_f

In order to apply our results on Fredholmness and the finite section method to $A = I + K$, we need to know about the limit operators of A , which, clearly, reduces to finding the limit operators of $K \in \mathcal{K}_f$. But before we start looking for these limit operators, we single out a subclass $\mathcal{K}_f^{\mathfrak{S}}$ of \mathcal{K}_f all elements of which are rich operators. So, this time, given $f \in \text{BUC}$, let

$$\begin{aligned} \mathcal{K}_f^{\mathfrak{S}} &:= \{ K \in \mathcal{K}_f : b_i \in \text{BUC}, c_i \in L_{\text{SC}\mathfrak{S}}^\infty \text{ for } i = 1, \dots, j \}, \\ \hat{\mathcal{B}}_{\mathfrak{S}} &:= \text{closspan}\{ M_b B M_c : b \in \text{BUC}, B \in R_f, c \in L_{\text{SC}\mathfrak{S}}^\infty \}, \\ \mathcal{B}_{\mathfrak{S}} &:= \text{closalg}\{ M_b B M_c : b \in \text{BUC}, B \in R_f, c \in L_{\text{SC}\mathfrak{S}}^\infty \}, \\ \hat{\mathcal{C}}_{\mathfrak{S}} &:= \text{closspan}\{ M_b C_a M_c : b \in \text{BUC}, a \in FL^1, c \in L_{\text{SC}\mathfrak{S}}^\infty \}, \\ \mathcal{C}_{\mathfrak{S}} &:= \text{closalg}\{ M_b C_a M_c : b \in \text{BUC}, a \in FL^1, c \in L_{\text{SC}\mathfrak{S}}^\infty \} \end{aligned}$$

denote the rich counterparts of $\mathcal{K}_f, \hat{\mathcal{B}}, \mathcal{B}, \hat{\mathcal{C}}$ and \mathcal{C} , and put

$$\mathcal{A}'_{\mathfrak{S}} := \text{closalg}\{ M_b, C_a M_c : b \in L_{\mathfrak{S}}^\infty, a \in FL^1, c \in L_{\text{SC}\mathfrak{S}}^\infty \}.$$

Recall that, by [25, Theorem 3.9], $BC \cap L_{\mathfrak{S}}^{\infty} = \text{BUC}$, and that $C_a M_c$ is rich for all $a \in FL^1$ and $c \in L_{\text{SC}\mathfrak{S}}^{\infty}$, whence every operator in $\mathcal{A}'_{\mathfrak{S}}$ is rich. Then the following “rich version” of Proposition 4.2 holds.

Proposition 4.3 *If $f \in \text{BUC}$, then it holds that*

$$\text{clos } \mathcal{K}_f^{\mathfrak{S}} = \hat{\mathcal{B}}_{\mathfrak{S}} = \hat{\mathcal{C}}_{\mathfrak{S}} = \mathcal{B}_{\mathfrak{S}} = \mathcal{C}_{\mathfrak{S}} \subset \mathcal{A}'_{\mathfrak{S}}.$$

In particular, every $K \in \mathcal{K}_f^{\mathfrak{S}}$ is rich.

Proof. All we have to check is that the arguments we made in the proof of Proposition 4.2 preserve membership of b and c in BUC and $L_{\text{SC}\mathfrak{S}}^{\infty}$, respectively. In only two of these arguments are there multiplications by b and c involved at all.

The first one is the proof of the inclusion $\hat{\mathcal{B}} \subset \hat{\mathcal{C}}$. In this argument, we show that every $B \in R_f$ is contained in $\hat{\mathcal{C}}$. But in fact, this construction even yields $B \in \hat{\mathcal{C}}_{\mathfrak{S}}$, which can be seen as follows. $B \in R_f$ is approximated in the operator norm by the operators $B^{(L)}$ from (25). Since the Courant hats φ_{ξ} and φ_{η} are in BUC and also $f \in \text{BUC}$, we get $\varphi_{\xi} \circ f \in \text{BUC}$ and $\varphi_{\eta} \circ f \in \text{BUC} \subset L_{\text{SC}\mathfrak{S}}^{\infty}$. So $B^{(L)} \in \hat{\mathcal{C}}_{\mathfrak{S}}$, and hence $B \in \hat{\mathcal{C}}_{\mathfrak{S}}$.

The second argument involving multiplication operators is the proof of the inclusion $\mathcal{C} \subset \hat{\mathcal{C}}$. But also at this point it is easily seen that the functions $b_m(x) = \int_{T_m} b(x-t) dt$ that are invoked in (26) are in fact in BUC , whence $\mathcal{C}_{\mathfrak{S}} \subset \hat{\mathcal{C}}_{\mathfrak{S}}$. ■

Now we are ready to say something about the limit operators of $K \in \mathcal{K}_f^{\mathfrak{S}}$. Not surprisingly, the key to these operators is the behaviour of the surface function f and of the multipliers b_i and c_i at infinity. We will show that every limit operator K_h of K is of the same form (2) but with f , b_i and c_i replaced by $f^{(h)}$, $b_i^{(h)}$ and $\tilde{c}_i^{(h)}$, respectively, in (3), where we use the notations introduced on page 11. We will even formulate and prove the analogous result for operators in $\mathcal{B}_{\mathfrak{S}}$. The key step to this result is the following lemma.

Lemma 4.4 *Let $B \in R_f$; that is, B is of the form (24) with a compactly supported kernel function $k \in C(\mathbb{R}^n \times [f_-, f_+]^2)$, and let $c \in L_{\text{SC}\mathfrak{S}}^{\infty}$. If a sequence $h = (h_m) \subset \mathbb{Z}^n$ tends to infinity and the functions $f^{(h)}$ and $\tilde{c}^{(h)}$ exist, then the limit operator $(BM_c)_h$ exists and is the integral operator*

$$\left((BM_c)_h u \right)(x) = \int_{\mathbb{R}^n} k(x-y, f^{(h)}(x), f^{(h)}(y)) \tilde{c}^{(h)}(y) u(y) dy, \quad x \in \mathbb{R}^n. \quad (28)$$

Proof. Choose $\ell > 0$ large enough that $k(r, s, t) = 0$ for all $r \in \mathbb{R}^n$ with $|r| \geq \ell$ and all $s, t \in [f_-, f_+]$. Now take a sequence $h = (h_m) \subset \mathbb{Z}^n$ such that the functions $f^{(h)}$ and $\tilde{c}^{(h)}$ exist. By definition of $f^{(h)}$ and $\tilde{c}^{(h)}$, this is equivalent to

$$\|f|_{h_m+U} - f^{(h)}|_U\|_{\infty} \rightarrow 0 \quad \text{and} \quad \|c|_{h_m+U} - \tilde{c}^{(h)}|_U\|_1 \rightarrow 0 \quad (29)$$

as $m \rightarrow \infty$ for every compact set $U \subset \mathbb{R}^n$. Moreover, it is easily seen that

$$(V_{-h_m} B M_c V_{h_m} u)(x) = \int_{\mathbb{R}^n} k(x-y, f(x+h_m), f(y+h_m)) c(y+h_m) u(y) dy$$

for all $x \in \mathbb{R}^n$ and $u \in \text{BC}$. Abbreviating $A_m := V_{-h_m} B M_c V_{h_m} - (B M_c)_h$, we get that $(A_m u)(x) = \int_{\mathbb{R}^n} d_m(x, y) u(y) dy$, where

$$\begin{aligned} & |d_m(x, y)| \\ &= \left| k(x-y, f(x+h_m), f(y+h_m)) c(y+h_m) \right. \\ &\quad \left. - k(x-y, f^{(h)}(x), f^{(h)}(y)) \tilde{c}^{(h)}(y) \right| \\ &\leq \left| k(x-y, f(x+h_m), f(y+h_m)) - k(x-y, f^{(h)}(x), f^{(h)}(y)) \right| \cdot \|c\|_\infty \\ &\quad + \|k\|_\infty \cdot \left| c(y+h_m) - \tilde{c}^{(h)}(y) \right| \end{aligned} \quad (30)$$

for all $x, y \in \mathbb{R}^n$ and $m \in \mathbb{N}$. Moreover, $d_m(x, y) = 0$ if $|x-y| \geq \ell$.

Now take an arbitrary $\tau > 0$, and denote by U and V the balls around the origin with radius $\tau + \ell$ and τ , respectively. Then, by (30),

$$\|P_\tau A_m\| = \text{ess sup}_{x \in V} \int_{\mathbb{R}^n} |d_m(x, y)| dy = \text{ess sup}_{x \in V} \int_U |d_m(x, y)| dy \rightarrow 0$$

as $m \rightarrow \infty$ since (29) holds and k is uniformly continuous. Analogously,

$$\|A_m P_\tau\| = \text{ess sup}_{x \in \mathbb{R}^n} \int_V |d_m(x, y)| dy = \text{ess sup}_{x \in U} \int_V |d_m(x, y)| dy \rightarrow 0$$

as $m \rightarrow \infty$. This proves that $(B M_c)_h$ from (28) is indeed the limit operator of $B M_c$ with respect to the sequence $h = (h_m)$. ■

Proposition 4.5 a) *Let $K = M_b B M_c$ with $b \in \text{BUC}$, $B \in R_f$ and $c \in L_{\text{SC}\mathfrak{S}}^\infty$. If $h = (h_m) \subset \mathbb{Z}^n$ tends to infinity and all functions $b^{(h)}$, $f^{(h)}$ and $\tilde{c}^{(h)}$ exist, then the limit operator K_h exists and is the integral operator*

$$(K_h u)(x) = \int_{\mathbb{R}^n} b^{(h)}(x) k(x-y, f^{(h)}(x), f^{(h)}(y)) \tilde{c}^{(h)}(y) u(y) dy, \quad x \in \mathbb{R}^n. \quad (31)$$

b) *Every limit operator of $K = M_b B M_c$ with $b \in \text{BUC}$, $B \in R_f$ and $c \in L_{\text{SC}\mathfrak{S}}^\infty$ is of this form (31).*

c) *The mapping $K \mapsto K_h$, acting on*

$$\{M_b B M_c : b \in \text{BUC}, B \in R_f, c \in L_{\text{SC}\mathfrak{S}}^\infty\},$$

as given in (31), extends to a continuous Banach algebra homomorphism on all of $\mathcal{B}_\mathfrak{S}$ by passing to an appropriate subsequence of h , if required. In particular, all limit operators K_h of $K \in \mathcal{K}_f \subset \mathcal{B}$ are of the form (2) with k replaced by

$$\hat{k}^{(h)}(x, y) = \sum_{i=1}^j b_i^{(h)}(x) k_i(x-y, f^{(h)}(x), f^{(h)}(y)) \tilde{c}_i^{(h)}(y). \quad (32)$$

Proof. **a)** From basic properties of limit operators [32, Proposition 1] we get that K_h exists and is equal to $(M_b)_h(BM_c)_h$ which is exactly (31) by Lemma 4.4.

b) Suppose $g \subset \mathbb{Z}^n$ is a sequence tending to infinity that leads to a limit operator K_g of K . Since $b, f \in L_{\mathfrak{S}}^\infty$ and $c \in L_{\text{SC}\mathfrak{S}}^\infty$, there is a subsequence h of g such that the functions $b^{(h)}$, $f^{(h)}$ and $\tilde{c}^{(h)}$ exist. But then we are in the situation of a), and the limit operator K_h of K exists and is equal to (31). Since h is a subsequence of g , we have $K_g = K_h$.

c) The extension to $\mathcal{B}_{\mathfrak{S}}$ follows from basic properties of limit operators [32, Proposition 1]. The formula for the limit operators of $K \in \mathcal{K}_f$ follows from the approximation of K by (27) for which we explicitly know the limit operators. ■

Example 4.6. Suppose $K \in \mathcal{K}_f$ where the surface function f and the functions b_i and c_i are all slowly oscillating. Let $h \subset \mathbb{Z}^n$ be a sequence tending to infinity such that $b_i^{(h)}$, $f^{(h)}$ and $\tilde{c}_i^{(h)}$ exist – otherwise pass to a subsequence of h with this property which is always possible.

From Example 3.3 we know that all of $b_i^{(h)}$, $f^{(h)}$ and $\tilde{c}_i^{(h)} = c_i^{(h)}$ are constant. Then, by Proposition 4.5 c), the limit operator K_h is the integral operator with kernel function

$$\hat{k}^{(h)}(x, y) = \sum_{i=1}^j b_i^{(h)} \tilde{c}_i^{(h)} k_i(x - y, f_h^{(h)}, f_h^{(h)}), \quad x, y \in \mathbb{R}^n \quad (33)$$

which is just a pure operator of convolution by $\hat{\kappa}^{(h)} \in L^1$ with

$$\hat{\kappa}^{(h)}(x - y) = \hat{k}^{(h)}(x, y) \quad (34)$$

for all $x, y \in \mathbb{R}^n$. □

5 The Main Results

The explicit formula (32) for the limit operators of K , together with our results on Fredholmness and the finite section method in terms of limit operators of A , gives us the desired criteria. These criteria are particularly explicit if all of the functions b_i , c_i and f are slowly oscillating, as in Example 4.6.

- In this case, Theorem 3.4 says that A is Fredholm if the Fourier transforms $F\hat{\kappa}_h$ of $\hat{\kappa}_h$ from (34) all stay away from the point -1 , and Theorem 3.5 says that this is a necessary condition for invertibility.
- Moreover, it will turn out that the BC-FSM is applicable to A if and only if A is invertible and all functions $F\hat{\kappa}_h$ stay away from -1 and have winding number zero with respect to -1 .

Here are the results in the more general case.

5.1 Fredholmness and invertibility

Let $f \in \text{BUC}$ and $K \in \mathcal{K}_f^{\mathfrak{S}}$. From Propositions 4.3 and 4.5 we know that $K \in \mathcal{A}_{\mathfrak{S}}'$, and all limit operators of $A = I + K$ are of the form $A_h = I + K_h$; that is

$$(A_h u)(x) = u(x) + \int_{\mathbb{R}^n} \sum_{i=1}^j b_i^{(h)}(x) k_i(x - y, f^{(h)}(x), f^{(h)}(y)) \tilde{c}_i^{(h)}(y) u(y) dy \quad (35)$$

for $u \in \text{BC}$ and $x \in \mathbb{R}^n$.

Now we apply Theorems 3.4 and 3.5 to our operator $A = I + K$.

Theorem 5.1 *If $f \in \text{BUC}$ and $K \in \mathcal{K}_f^{\mathfrak{S}}$, then*

$$\begin{aligned} A \text{ invertible on BC} &\Rightarrow \text{all limit operators (35) invertible on } L^\infty \\ &\Rightarrow A \text{ Fredholm on BC.} \end{aligned}$$

Proof. We just have to check that K is subject to all conditions in Theorem 3.4. Since, by Proposition 4.3, $\mathcal{K}_f^{\mathfrak{S}} \subset \mathcal{A}_{\mathfrak{S}}'$, K is rich. By Proposition 4.2, we have $K \in \mathcal{C}$. But since the generators of \mathcal{C} are band-dominated and the set of band-dominated operators is a Banach algebra, we get that all elements of \mathcal{C} , including K , are band-dominated. Moreover, every operator in \mathcal{C} is locally compact since L^1 -convolution operators are locally compact and multiplication operators commute with P_τ for all $\tau > 0$. ■

5.2 The BC-FSM

Since $K \in \mathcal{K}_f$ maps L^∞ into BC (see Remark 4.1 c)), we can, by Proposition 3.11, study the applicability of the BC-FSM (21) for $A = I + K$ by passing to its FSM (23) on L^∞ instead. This method is studied, for the case $n = 1$, in Theorem 4.2 in [24]. So let us restrict ourselves to operators on the axis, $n = 1$. By Theorem 4.2 in [24], we have to look at all operators of the form

$$QV_{-\tau}A_hV_\tau Q + P \quad \text{with} \quad A_h \in \sigma_+^{\text{op}}(A) \quad (36)$$

and

$$PV_{-\tau}A_hV_\tau P + Q \quad \text{with} \quad A_h \in \sigma_-^{\text{op}}(A) \quad (37)$$

with $\tau \in \mathbb{R}$, where $P = P_{[0, +\infty)}$ and $Q = I - P$. The operator (36) is invertible if and only if the operator that maps u to

$$u(x) + \int_{-\infty}^0 \sum_{i=1}^j b_i^{(h)}(x - \tau) k_i(x - y, f^{(h)}(x - \tau), f^{(h)}(y - \tau)) \tilde{c}_i^{(h)}(y - \tau) u(y) dy \quad (38)$$

with $x < 0$ is invertible on the negative half axis, or, equivalently,

$$u(x) + \int_{-\infty}^{\tau} \sum_{i=1}^j b_i^{(h)}(x) k_i(x-y, f^{(h)}(x), f^{(h)}(y)) \tilde{c}_i^{(h)}(y) u(y) dy, \quad x < \tau$$

is invertible on the half axis $(-\infty, \tau)$, for the corresponding sequence h leading to a limit operator at plus infinity.

And analogously, the operator (37) is invertible if and only if the operator that maps u to

$$u(x) + \int_0^{+\infty} \sum_{i=1}^j b_i^{(h)}(x-\tau) k_i(x-y, f^{(h)}(x-\tau), f^{(h)}(y-\tau)) \tilde{c}_i^{(h)}(y-\tau) u(y) dy \quad (39)$$

with $x > 0$ is invertible on the positive half axis, or, equivalently,

$$u(x) + \int_{\tau}^{+\infty} \sum_{i=1}^j b_i^{(h)}(x) k_i(x-y, f^{(h)}(x), f^{(h)}(y)) \tilde{c}_i^{(h)}(y) u(y) dy, \quad x > \tau$$

is invertible on the half axis $(\tau, +\infty)$, for the corresponding sequence h leading to a limit operator at minus infinity.

Theorem 5.2 *If $f \in \text{BUC}$ and $K \in \mathcal{K}_f^{\mathfrak{s}}$, then the BC-FSM is applicable to $A = I + K$ if and only if*

- A is invertible on L^∞ ,
- for every sequence h leading to a limit operator at plus infinity, the set $\{(38)\}_{\tau \in \mathbb{R}}$ is essentially invertible on $L^\infty(-\infty, 0)$, and
- for every sequence h leading to a limit operator at minus infinity, the set $\{(39)\}_{\tau \in \mathbb{R}}$ is essentially invertible on $L^\infty(0, +\infty)$.

Proof. Combine Proposition 3.11 above and Theorem 4.2 in [24]. ■

Remark 5.3. **a)** We say that a set $\{A_\tau\}_{\tau \in \mathbb{R}}$ is *uniformly invertible* if all A_τ are invertible and their inverses are uniformly bounded, and we call it *essentially invertible* if almost all A_τ are invertible and their inverses are uniformly bounded.

b) Both the operators (38) and (39) depend continuously on $\tau \in \mathbb{R}$. This implies that each ‘essentially invertible’ can be replaced by ‘uniformly invertible’ in the above theorem. We conjecture that, using the generalized collective compactness results of [13, 15], the words ‘essentially invertible’ can also be replaced by ‘elementwise invertible’ in Theorem 5.2.

c) If, as in Example 4.6, all of f , b_i and c_i are slowly oscillating, then we have $A_h = I + C_{F\hat{\kappa}^{(h)}}$ with $\hat{\kappa}^{(h)}$ as introduced in Example 4.6. In this case, by

Theorem 3.4, A is Fredholm if -1 is not in the spectrum of any $C_{F\hat{\kappa}^{(h)}}$; that is, all the (closed, connected) curves $F\hat{\kappa}^{(h)}(\mathbb{R}) \subset \mathbb{C}$ stay away from the point -1 . Moreover, the modified finite section method is applicable to A if and only if A is invertible and all curves $F\hat{\kappa}^{(h)}(\mathbb{R})$, in addition to staying away from -1 , have winding number zero with respect to this point.

d) In some cases (see Example 5.4 below) the functions $\hat{k}^{(h)}(x, y)$ from (33) in Example 4.6 even depend on $|x - y|$ only, which shows that the same is true for $\hat{\kappa}^{(h)}(x - y) := \hat{k}^{(h)}(x, y)$ then. If we then look at the applicability of the modified finite section method for $n = 1$, we get the following interesting result: The invertibility of A already implies the applicability of the finite section method. Indeed, if A is invertible, then, all limit operators A_h are invertible, which shows that all functions $F\hat{\kappa}_h$ stay away from the point -1 . But from $F\hat{\kappa}^{(h)}(z) = F\hat{\kappa}^{(h)}(-z)$ for all $z \in \mathbb{R}$ we get that the point $F\hat{\kappa}^{(h)}(z)$ traces the same curve (just in opposite directions) for $z < 0$ and for $z > 0$. So the winding number of the curve $F\hat{\kappa}^{(h)}(\mathbb{R})$ around -1 is automatically zero. \square

Example 5.4. Let us come back to Example 2.1 where, as we found out earlier, $n = 1$, $j = 2$, $b_1 \equiv -1/\pi$, $c_1 = f'$, $b_2 \equiv 1/\pi$, $c_2 \equiv 1$,

$$k_1(r, s, t) = \frac{r}{r^2 + (s - t)^2} - \frac{r}{r^2 + (s + t)^2}$$

and

$$k_2(r, s, t) = \frac{s - t}{r^2 + (s - t)^2} + \frac{s + t}{r^2 + (s + t)^2}.$$

In addition, suppose that $f'(x) \rightarrow 0$ as $x \rightarrow \infty$. Then, by Lemma 3.12 b) in [25], all of b_1, b_2, c_1, c_2 and f are slowly oscillating, and, for every sequence h leading to infinity such that the strict limit $f^{(h)}$ exists, we have that $b_1^{(h)} \equiv -1/\pi$, $c_1^{(h)} \equiv 0$, $b_2^{(h)} \equiv 1/\pi$, $c_2^{(h)} \equiv 1$, and $f^{(h)} \geq f_- > 0$ is a constant function, whence

$$\begin{aligned} \hat{k}^{(h)}(x, y) &= \frac{1}{\pi} \left(\frac{f^{(h)} - f^{(h)}}{(x - y)^2 + (f^{(h)} - f^{(h)})^2} + \frac{f^{(h)} + f^{(h)}}{(x - y)^2 + (f^{(h)} + f^{(h)})^2} \right) \\ &= \frac{2f^{(h)}}{\pi} \frac{1}{(x - y)^2 + 4(f^{(h)})^2} =: \hat{\kappa}^{(h)}(x - y), \quad x, y \in \mathbb{R}^n \end{aligned}$$

where $f^{(h)}$ is an accumulation value of f at infinity.

Now it remains to check the function values of the Fourier transform $F\hat{\kappa}^{(h)}$. A little exercise in contour integration shows that $F\hat{\kappa}^{(h)}(z) = \exp(-2f^{(h)}|z|)$ for $z \in \mathbb{R}$ (cf. Remark 5.3 d)). So $F\hat{\kappa}^{(h)}(\mathbb{R})$ stays away from -1 and has winding number zero.

Consequently, by our criteria derived earlier, we get that A is Fredholm and that the finite section method is applicable if and only if A is invertible.

As discussed in [31], by other, somewhat related arguments, it can, in fact, be shown that A is invertible, even when f is not slowly oscillating. Precisely, injectivity of A can be established via applications of the maximum principle

to the associated BVP, and then limit operator-type arguments can be used to establish surjectivity.

We note also that the modified version of the finite section method proposed in [10] could be applied in this case. (This method approximates the actual surface function f by an f for which f' is compactly supported before applying the finite section.) For this modified version the arguments of [10] and the invertibility of A establish applicability even when f is not slowly oscillating. \square

References

- [1] P. ANSELONE and I.H. SLOAN: Integral equations on the half-line, *J. Integral Eq.* **9** (1985), 3–23.
- [2] T. ARENS: Uniqueness for Elastic Wave Scattering by Rough Surfaces, *SIAM J. Math. Anal.* **33** (2001), 461–476.
- [3] T. ARENS: Existence of Solution in Elastic Wave Scattering by Unbounded Rough Surfaces, *Math. Meth. Appl. Sci.* **25** (2002), 507–528.
- [4] T. ARENS, S. N. CHANDLER-WILDE and K. HASELOH: Solvability and spectral properties of integral equations on the real line. II. L^p -spaces and applications, *J. Integral Equations Appl.* **15** (2003), 1–35.
- [5] K. E. ATKINSON: The numerical solution of integral equations on the half-line, *SIAM J. Numer. Anal.* **6** (1969), 375–397.
- [6] K. E. ATKINSON, *The numerical solution of integral equations of the second kind*, Cambridge University Press, 1997.
- [7] S. N. CHANDLER-WILDE, E. HEINEMEYER, AND R. POTTHAST: A well-posed integral equation formulation for 3D rough surface scattering, *Proc. R. Soc. Lond. A.*, published online 2006, DOI: 10.1098/rspa.2006.1752
- [8] S. N. CHANDLER-WILDE and C. R. ROSS: Scattering by rough surfaces: The Dirichlet problem for the Helmholtz equation in a non-locally perturbed half-plane, *Math. Meth. Appl. Sci.*, **19** (1996), 959–976.
- [9] S. N. CHANDLER-WILDE, C. R. ROSS and B. ZHANG: Scattering by infinite one-dimensional rough surfaces, *Proc. R. Soc. Lond. A.*, **455** (1999), 3767–3787.
- [10] S. N. CHANDLER-WILDE and A. MEIER: On the stability and convergence of the finite section method for integral equation formulations of rough surface scattering, *Math. Methods Appl. Sci.*, **24** (2001), 209–232.
- [11] S. N. CHANDLER-WILDE, M. RAHMAN and C. R. ROSS: A fast two-grid and finite section method for a class of integral equations on the real line with application to an acoustic scattering problem in the half-plane, *Numerische Mathematik*, **93** (2002), 1–51.

- [12] S. N. CHANDLER-WILDE and B. ZHANG: On the solvability of a class of second kind integral equations on unbounded domains, *J. Math. Anal. Appl.* **214** (1997), 482–502.
- [13] S. N. CHANDLER-WILDE and B. ZHANG: A generalised collectively compact operator theory with an application to second kind integral equations on unbounded domains, *J. Integral Equations Appl.* **14** (2002), 11–52.
- [14] S. N. CHANDLER-WILDE, S. LANGDON and L. RITTER: A high-wavenumber boundary-element method for an acoustic scattering problem, *Phil. Trans. R. Soc. Lond. A.*, **362** (2004), 647–671.
- [15] S. N. CHANDLER-WILDE and M. LINDNER: Generalized Collective Compactness and Limit Operators, submitted to *Integral Equations Operator Theory*.
- [16] B. DAHLBERG and C. KENIG: Hardy spaces and the Neumann problem in L^p for Laplace’s equation in Lipschitz domains, *Annals of Math.* **125** (1987), 437–466.
- [17] E. FABES, C. KENIG and G. VERCHOTA: The Dirichlet problem for the Stokes system on Lipschitz domains, *Duke Math. J.* **57** (1988), 769–793.
- [18] D. JERISON and C. KENIG: The Neumann problem on Lipschitz domains, *Bull. Amer. Math. Soc.* **4** (1981), 203–207.
- [19] K. JÖRGENS, *Linear Integral Operators*, Pitman, Boston, 1982.
- [20] C. E. KENIG, *Harmonic Analysis Techniques for Second Order Elliptic Boundary Value Problems*, American Mathematical Society, 1994.
- [21] M. LINDNER and B. SILBERMANN: Finite Sections in an Algebra of Convolution and Multiplication Operators on $L^\infty(\mathbb{R})$, *TU Chemnitz, Preprint* 6, 2000.
- [22] M. LINDNER and B. SILBERMANN: Invertibility at infinity of band-dominated operators in the space of essentially bounded functions, *In: Operator Theory – Advances and Applications* **147**, *The Erhard Meister Memorial Volume*, Birkhäuser 2003, 295–323.
- [23] M. LINDNER: Limit Operators and Applications on the Space of essentially bounded Functions, *PhD Dissertation, TU Chemnitz* 2003.
- [24] M. LINDNER: The Finite Section Method in the Space of Essentially Bounded Functions: An Approach Using Limit Operators., *Numerical Functional Analysis and Optimization* **24** (2003), 863–893.
- [25] M. LINDNER: Classes of Multiplication Operators and Their Limit Operators, *Journal for Analysis and its Applications* **23** (2004), 187–204.

- [26] M. LINDNER, *Infinite Matrices and Their Finite Sections: An Introduction to the Limit Operator Method*, Frontiers in Mathematics, Birkhäuser Basel, in press.
- [27] W. MCLEAN, *Strongly Elliptic Systems and Boundary Integral Equations*, Cambridge University Press, 2000.
- [28] Y. MEYER AND R. COIFMAN, *Wavelets: Calderón-Zygmund and Multilinear Operators*, Cambridge University Press, 1997.
- [29] M. MITREA: Generalized Dirac operators on nonsmooth manifolds and Maxwell's equations, *Journal of Fourier Analysis and its Applications* **7** (2001), 207–256.
- [30] D. MITREA, M. MITREA and M. TAYLOR: Layer potentials, the Hodge Laplacian, and global boundary problems in nonsmooth Riemannian manifolds, *Memoirs AMS* **150** no 713, 2001.
- [31] M. D. PRESTON, P. G. CHAMBERLAIN and S. N. CHANDLER-WILDE: An integral equation method for a boundary value problem arising in unsteady water wave problems, *In: Advances in Boundary Integral Methods: Proceedings of the 5th UK Conference on Boundary Integral Methods*, University of Liverpool 2005, 126–133.
- [32] V. S. RABINOVICH, S. ROCH and B. SILBERMANN: Fredholm Theory and Finite Section Method for Band-dominated operators, *Integral Equations Operator Theory* **30** (1998), 452–495.
- [33] V. S. RABINOVICH, S. ROCH and B. SILBERMANN, *Limit Operators and Their Applications in Operator Theory*, Operator Theory: Advances and Applications, **150**, Birkhäuser Basel 2004.
- [34] V. S. RABINOVICH and S. ROCH: The Fredholm index of locally compact band-dominated operators on $L^p(\mathbb{R})$, *TU Darmstadt, Preprint Nr. 2417*, 2005.
- [35] S. SAUTER and C. SCHWAB, *Randelementmethoden: Analyse, Numerik und Implementierung Schneller Algorithmen*, Teubner, 2004.
- [36] M. THOMAS: Analysis of Rough Surface Scattering Problems, *PhD Thesis, University of Reading*, in preparation.
- [37] R. H. TORRES and G. V. WELLAND: The Helmholtz equation and transmission problems with Lipschitz interfaces, *Indiana Univ. Math. J.* **42** (1003), 1457–1485.
- [38] G. C. VERCHOTA: Layer potentials and regularity for the Dirichlet problem for Laplace's equation in Lipschitz domains, *it J. Funct. Anal.*, **59** (1984), 572–611.

- [39] M. XIA, C. H. CHAN, S. LI, B. ZHANG and L. TSANG: An efficient algorithm for electromagnetic scattering from rough surfaces using a single integral equation and multilevel sparse-matrix canonical-grid method, *IEEE Trans. Ant. Prop.* **AP-51** (2003), 1142–1149.